

Biodiversity avoidance and wetland restoration in Cotonou, Benin

Modestine Victoire Bessan mbessan@worldbank.org¹; Randa Jean Nemer rnemer@worldbank.org¹; Tolidji Blaise Donou tdonou@worldbank.org¹; Van Anh Vu Hong vvuhong@worldbank.org²; Nahida Sinno² nsinno@worldbank.org; Alphonse Emadak aemadak@worldbank.org¹

1-Environmental Specialists, 2-Senior Urban Development Specialist, World Bank

Abstract

The latest of Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) report estimates that approximately 1 million animal and plant species are now threatened with extinction, an unprecedented event in human history. However, the report also indicates that this crisis is still reversible through immediate local and global actions aimed at preserving and restoring nature. In alignment with this call for local action to preserve ecological biodiversity, the Benin - Stormwater Management and Urban Resilience Project, funded by the World Bank, has implemented a biodiversity avoidance strategy as part of its water drainage infrastructure construction efforts. This paper details the approach and initial outcomes of the biodiversity avoidance and wetland restoration plan in the PA3 rainwater retention basin in Cotonou, Benin. Utilizing tablets and smartphones equipped with field navigation applications, along with drones for aerial imaging, this strategy has successfully preserved three enclaves of natural habitats covering an area of 9,420.55 m². These enclaves are home to numerous plant and animal species of significant socio-cultural, economic, and ecological importance. The next phase of this initiative will involve developing monitoring protocols for these biodiversity avoidance zones. By applying Artificial Intelligence (AI) and drone technology, the project will allow to accurately assess environmental changes in preserving habitats and optimize wildlife monitoring.

Keywords: Ecological biodiversity, wetlands, urban resilience, rainwater, Cotonou

1 Introduction

In the context of urban development, wetlands are particularly targeted and are destroyed to make way for urban infrastructure or solid sanitation task that completely change the structure of the wetlands. These wetlands are systematically targeted to house drainage and rainwater retention work. The construction of such work generally involves the destruction of these wetlands. The city of Cotonou, the economic capital of Benin, is the largest megalopolis in Benin with 1,228,667 inhabitants over an area of barely 68.5 km², i.e., a population density of 17,936.74 inhabitants/km². This port city of Cotonou is located in Ramsar 1018 site and therefore an international wetland. However, the city and nature are therefore generally considered incompatible, the city embodies the distance from nature. In urban areas, significant biological transformations appear that modify the ecology (Lougbeignon *and al.* 2011). Today, ecological concerns in urban areas are becoming increasingly increasing in view of the consequences that human activities generate on the health of city dwellers, especially in a city like Cotonou where compliance with environmental standards in terms of pollution and sanitation is a little weak. As a result, within the framework of the Cotonou City Stormwater Sanitation Program (PAPC) following several other urban projects such as the Urban Rehabilitation and Management Project (PRGU), the Urban Environmental Management Emergency Project (PUGEMU), etc., financed by Technical and Financial Partners (TFP), in this case the World Bank (WB), the requirement is increasingly made for compliance

with Environmental and Social Standards (ESS). In view of these very noble objectives, the thorny issue of nature arises, in this case the original habitats and ecosystems of the city of Cotonou through this ambitious project. Indeed, in the exposed and habitable parts where the city of Cotonou was built, the natural vegetation has almost entirely disappeared due to human actions. There are almost only alignment species, reforestation and very often exotic ornamental species. The current remnants of natural ecosystems of the city of Cotonou constitute the marshy areas of the city which are listed in the Ramsar 1018 site and currently constitute the natural retention basins to be developed and the water drainage outlets. For decision-makers and politicians, the primary interest of the urban populations of Cotonou lies in sanitation and the installation of water drainage work and therefore anything that can be "cumbersome" such as marsh vegetation must be cleared and removed, whereas for the ecological vision of a sustainable city in the objectives and ecological directives impose a share of "green" or natural to be saved in terms of ecological and sensitive habitat so that the city and its inhabitants continue to benefit from basic ecosystem services, especially in cities in developing countries such as Cotonou. It is imperative to establish within the framework of the PAPC an innovative approach that reconciles Environmental and Social Standards and urban sanitation measures decided by politicians. This justifies the development and implementation of a biodiversity avoidance plan during the construction of various drainage and water retention structures.

2 Material and Methods

2.1 Study area

2.1.1 Biophysical framework of the Pa3 rainwater retention basin in Cotonou

The Pa3 basin is located in the city of Cotonou in the 10th district. It is located between the parallels 6°22'55"N and 6°23'31" N and longitude 2°23'13' and 2°23'31" E. Figure 1 shows the geographical location map of the Pa3 basin. The Pa3 basin being in the city of Cotonou, benefits from a Guinean climate, better known locally as a subequatorial climate (Houeto, 2013). This climate is characterized by the annual succession of four seasons: 2 rainy seasons and 2 dry seasons alternating and of unequal durations (Bessan, 2013). Cotonou is one of the wettest cities in southern Benin with an average precipitation of 1346 ± 307 mm over the last thirty years (1990-2019) (Meteo-Benin, 2020). June is the wettest month with a rainfall of 329.30 mm. The average maximum monthly temperatures vary between 28.19° and 32.35°C at Cotonou station. In terms of hydrography, the Pa3 basin is part of the vast floodplain of Lake Nokoue.




Figure 1 : Geographical location of the Pa3 basin.

2.1.2 Socio-economic framework

The population living around the Pa3 basin is cosmopolitan, with several ethnic groups, practicing various beliefs. The activities practiced by the populations living near the Pa3 basin are mainly in the tertiary sector (trade, catering, administration, etc.) and the secondary sector (agri-food processing, mechanics, electricity, construction, crafts, etc.). Less than 1% of this population still practices market gardening and fishing activities. During rainy periods, these activities are disrupted by cyclical flooding around the Pa3 basin, making these populations vulnerable for a good part of the year.

2.2 Methodology for conducting work

2.2.1 Data collection equipment

The equipment used to carry out this work included tablets and smartphones equipped with the ODK Collect V1.30.1 and Locus Map 4.2 applications for navigation in the field and recording survey sheets; georeferencing in the field; fluorescent tape for demarcating avoidance zones; a digital camera for taking photos in the field; a pair of binoculars for bird watching; the analytical flora of Benin (Akoegninou *and al.*, 2006); a bird identification guide (Borrow & Demey, 2001); a mammal identification guide (Kingdon, 1997); a taxonomic identification guide for West African fish (Smith and al. 2009; Paugy *and al.*, 2003). Also, topographic and occupation maps of the Pa3 basin; flora and fauna survey sheets and a DJI Phantom 4 pro V2.0 drone for taking aerial images were used during the work. The use of drones will easily help to identify wildlife and monitor nesting or breeding areas of wildlife providing data without disturbing them. The photo 1 below shows the type of drone used.



Photo 1 : Phantom Pro 4 V2 used in this study (Photo taken by Lougbegnon 2023).

2.2.2 Methodological framework

A methodological framework session for the implementation of the strategy and protocol for the implementation of the biodiversity avoidance plan (flora and fauna) and the restoration of wetlands in the Pa3 rainwater retention basin was organized in the premises of the Urban Work Execution Agency with the presence of representatives of the PAPC steering team and Delegated Project Owner, the company in charge of the works and the control mission.

2.2.3 Communication and organization of a series of presentations on the importance of the identified avoidance zones, international wetlands and their importance and the need for resource conservation with various local stakeholders and company agents

After the ethnobiological surveys, various awareness-raising and communication phases on targeted themes were organized for both local stakeholders and local authorities and populations living in the Pa3 basin. These are the local elected officials from the districts which are concerned, the workers and technicians from the company in charge of the works and the Control Mission.

3 RESULTS

The various activities carried out as well as the technical devices have led to the achievement of the expected results in the Pa3 basin financed by the World Bank.

3.1 Identification and marking of natural habitats

The natural habitats to be spared have been identified and marked clearly and precisely in the Pa3 basin. This makes it possible to delimit these sensitive areas and guarantee their preservation (Photo 2).



Photo 2 : Partial view of the high *Thypha domengensis* meadow to be spared (E4) (above) and Pa3 basin avoidance zones delimited and marked (bottom) (Houessou Laurent, 2023).

3.2 Mapping of natural habitats and area of avoidance zones

A total of three avoidance zones were mapped, with a total area of 9420.55 m². These areas, named E2, E4 and E5, were clearly demarcated, marked and identified to ensure that they are not disturbed during the work. The mapping of the natural habitats to be spared has been successfully completed. The location and characteristics of these habitats are now available as map data. In addition, the land use in the Pa3 retention basin has also been mapped.




Figure 2 : Final mapping of natural habitats (on the left) and that of occupations (on the right) to be spared on the Pa3 retention basin.

3.3 Identification and protection of important species

Species of great socio-cultural and ecological importance were identified and protected. Measures were taken to delimit and materialize the avoidance zones using fluorescent strips and indicative plates. Several plant species are found in these ecosystems and are essential for documenting the biodiversity of the Pa3 basin and are useful for conservation. In particular, the conservation of *Thalia geniculata*, a plant species used as packaging leaves (Plate 1) and having cultural and economic values.



Plate 1 : Collection and piling of *Thalia geniculata* leaves for marketing (Photo taken by Houessou, 2023).

The Pa3 basin is home to a remarkable diversity of birds typical of the lake, lagoon and river environments of southern Benin. These birds are often associated with aquatic habitats and play an important role in the local ecosystem by feeding mainly on fish and other aquatic organisms. The count of 17 bird species around the Pa3 basin is evidence of the richness of the avian fauna of the basin's wetlands. This diversity of bird species can be an indicator of the health of the aquatic ecosystem and the quality of the habitat. It is essential to monitor and preserve the enclaves to ensure the conservation of these avian species.

3.4 Information/awareness/training sessions for the various stakeholders in the Pa3 retention basin

This session took into account technicians, machine operators, site managers, team leaders, workers from the company in charge of the work and the control mission whose objective is to prepare these various stakeholders to: (i) know the three enclaves of the Pa3 basin delimited as avoidance zones; (ii) raise awareness among these various field stakeholders on the precautions to take to safeguard and protect the three avoidance zones delimited during the work.

3.5 Raising awareness among the population on measures to avoid probable incidents related to the dredged areas

In search of fish and other resources, the population, especially young people and children, often try to enter the dredged portions. To avoid any possible damage from such practices during the work, the populations surrounding the basin were made aware by the team of experts and the company of the dangers to which they may be subjected when entering the site and going near the open drain for any activity. Plate 2 presents the situation of this event and the presence of an awareness team.

3.6 Barricade and installation of signage in the access areas to the site

In addition to raising awareness, other measures were taken. These measures include the installation of barricades and the installation of danger signage to prevent access to the sites by the populations. The team of experts accompanied the company team in the installation of these various signage structures. Plate 2 shows some images of the installation of prohibition signs.



Plate 2 : Installation of signalization panels along the Pa3 basin (Photo taken by Lougbegnon, 2023).

4 CONCLUSION AND RECOMMENDATIONS

This article presents a summary of the methodology adopted and some salient results of the activities carried out during the implementation of the strategy and protocol of the biodiversity avoidance plan and the restoration of wetlands in the rainwater retention basin Pa3 of Cotonou. This is a new concept introduced as part of the PAPC sanitation tasks in Benin. As part of the avoidance mission, it should be noted that most of the activities carried out are focused on the delimitation of the avoidance zones, their securing and awareness-raising activities. Thus, three enclaves of natural habitats to be spared have been clearly delimited and marked. These enclaves cover an area of 9420.55 m² with well-defined characteristics. A distribution map of these enclaves has been produced. Added to this, there are workshops and awareness-raising sessions for key stakeholders who should help leading the process. In view of these different activities carried out using new technologies; natural habitats are identified and clearly marked in the basins financed by the World Bank and the avoidance zones are marked and monitored during the tasks, the mapping of natural habitats to be spared is carried out and available and species of great socio-cultural and ecological importance are identified and protected. However, it will be interesting in the short and medium term to develop biodiversity monitoring protocols around this Pa3 basin in order to properly assess i-) environmental changes in the avoidance zones; ii-) optimal monitoring of wildlife. The use of drones will easily help to identify wildlife and monitor nesting or breeding areas of wildlife providing data without disturbing them. Given that these basins are migration sites for Palearctic or intra-African migratory birds, the use of drones will also make it possible to monitor and make rapid counts of birds in these basins after the work to know the rates of colonization or reconstitution of ecosystems because birds are good indicators for this purpose.

5 REFERENCES

- Akoegninou, A., van der Burg, W. J., & van der Maesen, L. J. G. (Eds.) (2006).** Analytical flora of Benin. (Wageningen Agricultural University papers; No. 06.2). Backhuys Publishers. <https://edepot.wur.nl/281595>
- Bessan M. V. (2013):** Evaluation of public policies for the management of natural risks and disasters in Benin. End of study work for the Master of Sciences “Management of Natural Disaster and Risk”. University of Liège in Belgium (ULg)/Faculty of Sciences-Institute of Geography, 85 pages.
- Borrow N & Demey R. (2001).** A Guide to the Birds of Western Africa – Hardcover. Published by Princeton University Press, c. 2001, 2001.
- Houeto O. F., (2013).** Mapping of sacred forests in Ramsar sites 1017 and 1018 in Benin. Report. ITTO, Yokohama, 81 p.
- IPBES 11 (2024)** Summary of the 11th Session of the Plenary of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. In Earth Negotiations Bulletin, Vol. 31 No. 77, Thursday, 19 December 2024. <https://enb.iisd.org/sites/default/files/2024-12/enb3177e.pdf>
- IPBES (2019):** Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. S. Díaz, J. Settele, E. S. Brondízio E.S., H. T. Ngo, M. Guèze, J. Agard, A. Arneth, P. Balvanera, K. A. Brauman, S. H. M. Butchart, K. M. A. Chan, L. A. Garibaldi, K. Ichii, J. Liu, S. M. Subramanian, G. F. Midgley, P. Miloslavich, Z. Molnár, D. Obura, A. Pfaff, S. Polasky, A. Purvis, J. Razzaque, B. Reyers, R. Roy Chowdhury, Y. J. Shin, I. J. Visseren-Hamakers, K. J. Willis, and C. N. Zayas (eds.). IPBES secretariat, Bonn, Germany. 56 pages. https://files.ipbes.net/ipbes-web-prod-public-files/2020/02/ipbes_global_assessment_report_summary_for_policymakers_fr.pdf
- Kingdon, J. (1997)** The Kingdon Field Guide to African Mammals. Academic Press, London.
- Loughbenon, J. Codjia, R. Libois (2011).** Distribution of avifauna in alternative forest environments (plantations and fallows) in southern Benin in relation to habitat factors. Published 7 January 2011. Geography. International Journal of Biological and Chemical Sciences
- Meteo-Benin, (2020).** Provision of rainfall and temperature data for the city of Cotonou from 1990 to 2019)
- Paugy D., Lévêque C. & Teugels G. G. (2003).** — The Fresh and Brackish Water Fishes of West Africa, vols. 1 & 2. National Museum of Natural History, Paris, 1272p. (Tropical Fauna and Flora; 40).
- Profizi J-P., Ardila-Chauvet S., Billot C., Couteron P., Delmas M., Diep T. M. H., Grandcolas P., Kokou K., Muller S., Rana A. S., Ranarijaona H. L. T., and Sonke B. (2022).** Biodiversity of Intertropical Ecosystems: Knowledge, Sustainable Management, and Valorization. IRD Éditions. Marseille, 782 pages
- Smith, K.G., Diop, M.D., Niane, M. and Darwall, W.R.T. (Compilers). (2009).** The Status and Distribution of Freshwater Biodiversity in Western Africa Gland, Switzerland and Cambridge, UK: IUCN. x+94pp+4pp cover. ISBN: 978-2-8317-1163-8